



BLACK WALNUT TOXICITY

Background

Walnut trees are all part of the genus *Juglans*. There are some 15 species of deciduous walnut trees, sometimes shrubs, found in S.E. Europe, Asia, North and South America. Some of these species include *J. californica* (Californian walnut), *J. cinerea* (Butternut), *J. regia* (English or Persian walnut), *J. nigra* (Black Walnut), and *J. sieboldiana* (Japanese walnut). Walnuts are grown for their fruit (what is ordinarily called a nut). In addition, some species are grown for the dark wood used to construct furniture.

J. nigra (Black Walnut) is a long-lived, vigorous spreading tree with pinnate, aromatic leaves up to 24-inches long, each with 11 to 23 oblong, glossy, dark green leaflets. This tree can grow to over 100-feet tall, attaining a spread of up to 70 feet.

Problem

The Black Walnut (*J. nigra*) produces a chemical called juglone that is toxic to many plant species. Highest concentrations of this chemical are found in buds, leaves, nut husks and roots. The chemical is not very soluble, but it accumulates under the tree's drip zone. (The drip zone is the area on the ground measured from the trunk of a tree or shrub to the outer edge of its branch reach.) Typically the chemical is released by decaying leaves and roots. Death or poor growth of other plants may occur under the spread of Black Walnut trees due this chemical. Other species of the *Juglans* produce smaller amounts of juglone.

Plants sensitive to juglone include alfalfa, apple, birch, blackberry, blueberry, cabbage, eggplant, Kalmia, lilac, magnolia, peony, pepper, potato, pine, privet, rhododendron, some rose varieties, silver maple, and tomato.

Resistant plants to juglone include beans, beets, cedar, cherry, corn, daffodil, elm, ferns, forsythia, hawthorn, hemlock, iris, locust, most maples, oaks, onions, orchard grass, pachysandra, sycamore, viburnums, and wheat.

Susceptible plants grow poorly or not at all next to or under Black Walnut trees. Stems of these plants may show blisters and blighted spots where Black Walnut leaves are used as winter protective cover.

Continued on back page

Control

1. Consider neighboring plants, soil pH, moisture, drainage, and exposure to sun and wind when selecting a Black Walnut for your property.
2. Locate sensitive plants well away from all walnut trees.
3. Avoid accumulation of leaves and nut hulls under and around the trees. Raked debris should be stored and composted separately.
4. Do not use walnut leaves or hulls as mulch around susceptible plants.

References

Peattie, D.C., A Natural History of Western Trees. Bonanza Books, N.Y.

2002 PNW Plant Disease Management Handbook

Plant Encyclopedia

Hortis Third – Dictionary

10,000 Garden Questions Answered

Gray, A., 1887, Gray's Lessons and Manual of Botany. American Book Company.

Sustainable Gardening. The Oregon-Washington Master Gardener Handbook. WSU Extension
